

HEADLICE POLICY

Inspirational Learning Academies Trust is aware of the national problem of head lice and how it can sometimes affect children of a primary school age at home and in school.

This policy attempts to set out the duties and responsibilities of parents, the school and the Health Authorities in dealing with head lice. It sets out what school can and cannot do legally as well as providing some practical advice as to how to tackle head lice.

Parent's/Carers/Guardians Duties

As the carer of a child you should;

- Undertake regular weekly and where appropriate daily checks to ensure that your child/family is not affected. This should be done with a detector comb as noted in the advice section. All family members should be checked.
- Seek advice from a medical practitioner (Pharmacist/ Health Visitor/School Nurse/Doctor) about using an approved lotion and treat using this method in the recommended way OR Use 'Wet Combing Method'

School will undertake to

- Provide information about Headlice on an annual basis and more regularly if many cases arise.
- Encourage parents to seek advice from the School Nursing Service if they find Head lice to be a problem.
- Maintain a visual check on pupils.
- Alert parents (class/year group) via a text/ letter if an outbreak is noted
- Alert individual parents by letter/ phone call if a more persistent case is noted.
- Invite these parents to meet with the Safeguarding Officer to discuss treatment/ provide advice.
- The Safeguarding Officer (with the parent's permission) will then; refer that parent/child to the School Nurse/ Health Visitor enabling that parent to gain support.
- Offer to provide assistance with wet combing and /or other treatment.

The school by law cannot

- Carry out physical checks on pupils for head lice.
- Tell parents to keep children away from school because of head lice.
- Exclude a child from school because of head lice.

HEADLICE POLICY

Advice Section

Facts about head lice

- Head lice are small, six-legged wingless insects that live on the hair close to the scalp where it is warm and easy for them to feed.
- They are pin-head size when they hatch, less than match-head size when fully grown and are difficult to see in the hair.
- The eggs are glued individually to hairs near the scalp. Unhatched eggs are dull in colour and hard to see, but after the lice have hatched the empty egg sacs - called 'nits' - are white and easy to see. Many people mistake the empty egg sacs or nits for head lice when they are actually evidence of a previous infection of head lice.
- Head lice are easily missed in dry hair and do not necessarily cause itching. There are often only 10 lice or less on a head.
- Head lice can't fly, jump or swim. They spread from person to person by climbing swiftly along hairs during close head to head contact.
- Head lice are not fussy about hair length or condition - clean hair is no protection.
- Head lice are not harmful.
- Head lice infection is common. Anyone with hair can get them, but children, who put their heads together a lot, tend to get them more often.

Detection

- Head lice can't be prevented but regular checking ensures early detection and treatment if necessary. The best detection method is wet combing (see below).
- Parents and carers should aim to check their children's hair once a week during hair washing. You need your usual shampoo, ordinary conditioner and a louse detection comb. Remember that you are looking for living moving head lice - the only evidence that your child has a head lice infection. The comb must be fine enough to catch the lice. Your pharmacist should be able to recommend a suitable one.

How to check

- Good lighting is important. Look for nits by parting hair in small sections, going from one side of the head to the other. Check carefully, looking close to the scalp.
- Nits are most predictably found on hairs at the nape of the neck and behind the ears, where they are protected from extremes of light and temperature. However, they may be laid anywhere on the hair, especially in warm weather.
- The appearance of a nit is often confused with that of a flake of dandruff or a dried particle of hairspray or gel. A distinguishing feature is that dandruff and hair products can be easily combed off the hair or removed with the fingers, while nits cannot. Nits are firmly glued to the hair and must be removed with a fine toothed comb or fingernails, or snipped off with scissors.
- The scalp should be examined in sunlight or under bright artificial light. The hair should be parted, with individual strands checked for nits. Head lice 'hold on' to the hair with hook like claws found at the end of each of their six legs.

HEADLICE POLICY

Treatment

There are two treatment options. The first is the wet combing method as set out below. Wet combing is by far the most effective method. The other is to use an insecticide lotion.

"Persistence is the key to getting rid of these annoying insects!"

Wet combing

1. Shampoo the hair, rinse, apply lots of conditioner and use a wide-toothed comb to untangle and straighten.
2. Once the tangles have gone, section the hair and comb with a fine-toothed detection comb from root to end. Comb one section at a time, paying particular attention to the back of the neck and the area around the ears.
3. Check the comb for live lice between each stroke and remove them.
4. When the whole head of hair has been combed through and checked, rinse off the conditioner.
5. Check all family members at the same time and arrange treatment if lice are found.
6. If you discover head lice, inform any close contacts, so they and their families know they should check too.
7. The time required to wet comb the whole head can range from 2 minutes for short, straight hair to 30 minutes for long, curly hair.
8. The wet combing method needs to be repeated every 3 days for up to 3 weeks to ensure that all head lice are removed.

Lotions

- You can ask your school nurse, health visitor, pharmacist or doctor about which lotion is most effective. People with asthma or allergies, pregnant or breastfeeding mothers, and parents or carers of very young children should always ask their doctor or pharmacist about which treatments are safe for them.
- Note the following advice for using lotions:
- Do not use lotions unless you find a living, moving louse. Head lice may become resistant to lotions if they are used too often.
- If live head lice are still being found after two treatments, ask your doctor, pharmacist, health visitor or school nurse for advice.
- Follow the instructions on the packet carefully. These will tell you how to apply the lotion, how long to leave it on the hair and when any repeat application is required for the treatment to be effective. The treatment method may vary between different lotions.
- Although the lotion will kill the lice, it will not kill the eggs. You will need to apply the lotion again after 7 days to kill any newly hatched baby lice. Again, this procedure may vary between different lotions, so follow the instructions on the packet carefully.

HEADLICE POLICY